BREAST AUGMENTATION

DR. GERUT'S PREOPERATIVE INSTRUCTIONS

FOR OFFICE SURGERY
Patients are required to have preoperative blood tests at their doctor's office or at the laboratory of their choice. The tests should be done no more than ten days before surgery and the results must reach the office no less than two days before surgery.
EOD HOCDITAL CUDCEDY
Patients must visit the hospital BEFORE the day of surgery for pre-operative blood tests and interviews with an anesthesiologist and a nurse. Some hospitals allow the blood tests to be done at your doctor's office (if the doctor's lab is state approved) or at a private laboratory. Even if your tests are done outside the hospital, you must have a presurgical interview at the hospital.
A MEDICAL EXAMINATION and EKG may be required before surgery. Please understand that if this is required, it is for your benefit and safety.
DO NOT TAKE ASPIRIN, BABY ASPIRIN, BUFFERIN, APC, FIORINAL, ALKA SELTZER, ECOTRIN, EXCEDRIN, ANACIN, VITAMIN E, VITAMIN C or any other medication for two weeks before surgery except as prescribed by Dr. Gerut. Even Advil, Motrin, Aleve, etc., should be avoided. If necessary you may take Tylenol. Please check with the office if there are any questions regarding medication.
NO SMOKING for at least two weeks prior to surgery as it drastically affects healing and lung function. It can cause the implants to build up scar tissue and move or feel hard as well as other problems.
Not more than one alcoholic drink per day as of one week before surgery as alcohol can cause excessive swelling.
PRE-OP MEDICATION To be taken BEFORE surgery (only the items indicated by check mark):
Zofran - Anti-nausea. Take pill, with a tiny sip of water, 3 hours before surgery.
☐ Vitamin K1 Twice a day for 5 days before surgery (not morning of surgery).
Hibiclens soap Wash from shoulders to belly button twice a day for 5 days before surgery and including morning of surgery
☐ Multivitamin 1 daily
Please stop ALL natural or holistic medications as they may interfere with your prescribed medications and anesthesia. The one exception is arnica in moderate doses.
Prior to surgery you should have a MAMMOGRAM / SONOGRAM depending on your age.
<u>Nothing</u> to eat or drink after midnight the night before surgery. <u>Nothing</u> to eat or drink the morning of your surgery except the Emend with a sip of water.

Please be sure you have a thermometer at home. You must take your temperature daily for a week after surgery.

Please wear loose fitting clothing that buttons or zippers down the front and low shoes or sneakers.

Please do not wear make-up, perfume, face moisturizer, or body moisturizer on the day of surgery.

Please tell us if you have ever had any skin conditions such as cold sores, shingles, herpes, etc.

Please tell us if you have ever taken Accutane, blood thinner, birth contol pills, thyroid medicine or any hormone pills.

Arrange for someone to pick you up at the hospital or office after surgery and to stay with you at home for at least 24 hours. Please bring their phone number with you.

If you develop symptoms of illness such as a cold or a cough, please call our office immediately at 516-295-2100.

DO NOT SHAVE UNDERARMS THE DAY OR NIGHT BEFORE SURGERY. YOU MAY SHAVE TWO DAYS BEFORE SURGERY OR THE MORNING OF SURGERY.

I have been made aware of the following possible complications and/or side effects and I accept them (there are others possible that are not listed here):

Reaction to the sutures

Bleeding	Thrombophlebitis	Reaction to the sutures
Infection	Hypertrophic Scars	Capsule Formation (hardening)
Need for revisional surgery	Keloids	Skin/scar discoloration
Wound Dehiscence	Chronic Pain	Disability
Skin Numbness	Asymmetry – very likely	Change in breast shape
Seroma	Anesthesia Risks	Paresthesias
Loss of nipple areolar sensation	Nipple areolar malposition	Scarring
Change in Mammogram	Hematoma	Contour deformities
Change of breast shape with che	st muscle motion	Spontaneous deflation of implant
Implant can leak and needs immediate replacement		Interference with mammogram
Need to replace implant after 10	years (silicone) even with no leakage	Change in breast self-examination
	X	
If you wish to take any sedatives or	tcal condition between now and the day of your surgery your consend possible complications, my questions	_
Dationt Cionatura	Doto	
Patient Signature	Date	
Zachary E. Gerut, M.D., F.A.C.S., P.C. Diplomat, American Board of Plastic Surger		

Diplomat, American Board of Plastic Surgery
Member, American Society of Plastic Surgeons
Member, American Society for Aesthetic Plastic Surgery
Assistant Clinical Professor of Plastic Surgery, Albert Einstein Med Ctr. (ret)
Medical Director, Touro University Physician Assistant School
Fellow of the American College of Surgeons