FAT INJECTION

DR. GERUT'S POST OPERATIVE INSTRUCTIONS

Avoid, talking, chewing, etc. Do not use straws. DO NOT CHEW GUM OR SMOKE!

On the first night after surgery, avoid excessive chewing. After the second day you may eat or drink anything that you wish.

Please take the medications that Dr. Gerut indicates with a check mark. Only the medications checked off will be given to you. If you are allergic to any of these medications, please let us know! Please take all pills with a bit of food to avoid nausea. When you fill your prescriptions, the bottles will be labeled only with the medical names for your pills and for your reference, these medical names are in parentheses below.

Antibiotic:	□ Keflex (Cephalexin) 500mg (1pill) 2 times a day Or, if allergic to Penicillin,		
	□Vibramycin (Doxycycline) - 100mg (1pill) 2 times a day		
Severe pain:	□Percocet (Oxycodone) - 1-2 pills every 3 to 4 hours as needed.		
Moderate pain:	■ Norco (hydrocodone bitartrate/acetaminophen)- 1-2 pills every 3 to 4 hours as needed.		
Nausea:	☐ Compazine (Prochloperazine) Use one if needed. Never use more than 2 in 24 hours. If nausea returns, and you weigh less than 160 pounds, call the office before using a second suppository.		
Constipation:	□Ex-Lax (or any other mild laxative as needed).		
Multi-Vitamin:	□1 daily		

There will be a certain amount of swelling that will last a variable number of days. To minimize swelling, keep your head elevated at all times. When lying down or sleeping, sleep in a reclining chair or use as may pillows as possible. You may apply ice if you wish. If you have a any problem especially a sudden increase in pain or a fever (measured by a thermometer) or if you have MUCH more swelling on one side compared to the other, **call the office at 516-295-2100**. After office hours please call **516-620-3619**.

I have been made aware of the followhem (there may be other problems		ons and/or side effects and I accept
Bleeding Need for revisional surgery Change in facial shape Parasthesias and/or numbness Change in facial expression X	Reaction to the sutures Chronic pain Seroma Hematoma	Infection Asymmetry Anesthesia risks Contour deformities
over the Internet from which you macknowledge and accept this poten	involve sending informating or may not be identifiatial breach of your privacy	on about or photographs of yourself ble. By your initials here you . X
No Aspirin, Bufferin, APC, Fiorina other medication not directed by D Advil or Motrin, check with Dr. Genarcotic pain reliever, you are to avsevere drowsiness can cause falls of	r. Gerut for ONE WEEK A erut or his office staff. Dur oid driving or operating he	AFTER SURGERY. Regarding ring the time that you are taking a
Significant weight gain or weight leasurgical result, For this reason, you after your procedure. If your weight weight) after your procedure, you was area and you will be charged fee. If for other reasons, revisional there is no surgical fee, THERE WAROOM AND THE ANESTHESIA	must agree to make all effort changes significantly (clayill not be entitled to any formal a surgical fee and an operation of the surgery is required, there results a STANDARD F.	forts to keep your weight stable hange of 10% or more in body ree "touch-up" procedures to the ating room fee and an anesthesia may be a surgical fee but even if
I have read the above instructions (have been answered and I understa	1 0	
Patient Signature		Date
Zachary E. Gerut, M.D., F.A.C.S., P.C. Diplomat, American Board of Plastic Surg Member, American Society of Plastic Surg Member, American Society for Aesthetic Assistant Clinical Professor of Plastic Surg	geons Plastic Surgery	ret)

rev 6/16

Medical Director or Research, Touro University Physician Assistant School

Fellow of the American College of Surgeons